

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: *J Allergy Clin Immunol* 1996 Jan;97(1 Pt 1):53-64 Related Articles, Books, LinkOutPubMed
Services

Single amino acid substitutions on a Japanese cedar pollen allergen (Cry j 1)-derived peptide induced alterations in human T cell responses and T cell receptor antagonism.

Ikagawa S, Matsushita S, Chen YZ, Ishikawa T, Nishimura Y


Department of Neuroscience and Immunology, Kumamoto University Graduate School of Medical Sciences, Japan.

Related
Resources

We generated T cell clones specific to a Japanese cedar pollen allergen (Cry j 1) and investigated effects of altered T cell receptor (TCR) ligand on changes of T cell responses. One of these Cry j 1-specific T cell clones established from patients with Japanese cedar pollinosis, ST1.9, recognized an antigenic peptide Cry j 1 p335-346 in the context of HLA-DRA+DRB3*0301 molecules and secreted interleukin-4 dominantly, with a smaller amount of interferon-gamma. ST1.9 represented one of the major T cell clones specific to Cry j 1 in the donor, because a short-term cultured polyclonal T cell line specific to Cry j 1 exhibited the same character as the ST1.9. We synthesized various analog peptides derived from Cry j 1 p335-346 with single amino acid substitutions and determined key residues for interactions between TCR of ST1.9 and HLA-DR molecules. We also analyzed changes in the responses of ST1.9 to Cry j 1 p335-346-derived analog peptides. Of interest was that the substitution of 339threonine to valine resulted in a significant increase in interferon-gamma production, with no remarkable changes either in proliferative response or interleukin-4 production. Analog peptides carrying the substitutions of 339threonine to glycine or glutamine revealed TCR antagonism, without changes in their binding affinities to the DR molecule. Therefore single amino acid substitutions on an allergen peptide carrying the T cell epitope may suppress helper-T-dependent class switch pressure to IgE in B cells either by inducing increased interferon-gamma production or by inhibiting proliferative responses in helper-T cells.

PMID: 8568138

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	
Search PubMed	▼	for					Go	Clear
Limits Preview/Index History Clipboard								

Entrez PubMed

Display	Summary	▼	Save	Text	Order	Add to Clipboard
Show: 20	▼	Items 1-20 of 216	Page 1 of 11		Select page: 1 2 3 4 5 6 7 8 9 10 >>	

PubMed
Services

- ☐ 1: [Sone T, Morikubo K, Miyahara M, Komiyama N, Shimizu K, Tsunoo H, Kino K.](#) Related Articles
T cell epitopes in Japanese cedar (*Cryptomeria japonica*) pollen allergens: choice of major T cell epitopes in Cry j 1 and Cry j 2 toward design of the peptide-based immunotherapeutics for the management of Japanese cedar pollinosis.
J Immunol. 1998 Jul 1;161(1):448-57.
PMID: 9647255


Related
Resources

- ☒ 2: [Hashiguchi S, Sugimura K.](#) Related Articles
[Molecular immunology of Japanese cedar pollen allergens: analysis of T cell epitopes].
Nippon Rinsho. 1996 Aug;54(8):2233-42. Review. Japanese.
PMID: 8810803
- ☐ 3: [Hori T, Kamikawaji N, Kimura A, Sone T, Komiyama N, Komiyama S, Sasazuki T.](#) Related Articles
Japanese cedar pollinosis and HLA-DP5.
Tissue Antigens. 1996 Jun;47(6):485-91.
PMID: 8813737
- ☐ 4: [Sone T, Morikubo K, Shimizu K, Komiyama N, Tsunoo H, Kino K.](#) Related Articles
Peptide specificity, HLA class II restriction, and T-cell subsets of the T-cell clones specific to either Cry j 1 or Cry j 2, the major allergens of Japanese cedar (*Cryptomeria japonica*) pollen.
Int Arch Allergy Immunol. 1999 Jul;119(3):185-96.
PMID: 10436390
- ☐ 5: [Ohno N, Ide T, Sakaguchi M, Inouye S, Saito S.](#) Related Articles
Common antigenicity between Japanese cedar (*Cryptomeria japonica*) pollen and Japanese cypress (*Chamaecyparis obtusa*) pollen, II. Determination of the cross-reacting T-cell epitope of cry j 1 and cha o 1 in mice.
Immunology. 2000 Apr;99(4):630-4.
PMID: 10792512
- ☐ 6: [de Lalla C, Sturniolo T, Abbruzzese L, Hammer J, Sidoli A, Sinigaglia F, Panina-Bordignon P.](#) Related Articles
Cutting edge: identification of novel T cell epitopes in Lol p5a by computational prediction.
J Immunol. 1999 Aug 15;163(4):1725-9.
PMID: 10438899

- ☐ 7: [Abe Y, Kimura S, Kokubo T, Mizumoto K, Uehara M, Katagiri M.](#) Related Articles
Epitope analysis of birch pollen allergen in Japanese subjects.
J Clin Immunol. 1997 Nov;17(6):485-93.
PMID: 9418189
- ☐ 8: [Muller WD, Karamfilov T, Bufe A, Fahlbush B, Wolf I, Jager L.](#) Related Articles
Group 5 allergens of timothy grass (Phl p 5) bear cross-reacting T cell epitopes with group 1 allergens of rye grass (Lol p 1).
Int Arch Allergy Immunol. 1996 Apr;109(4):352-5.
PMID: 8634519
- ☐ 9: [Ishikawa T, Ikagawa S, Masuyama K, Matsushita S, Nishimura Y.](#) Related Articles
Human T cell response to antigen peptides of Japanese cedar pollen (Cry j 1).
Int Arch Allergy Immunol. 1997 May-Jul;113(1-3):255-7. No abstract available.
PMID: 9130539
- ☐ 10: [Wurtzen PA, Bufe A, Wissenbach M, Madsen HO, Ipsen H, Arned J, Van Neerven RJ.](#) Related Articles
Identification of isoform-specific T-cell epitopes in the major timothy grass pollen allergen, Phl p 5.
Clin Exp Allergy. 1999 Dec;29(12):1614-25.
PMID: 10594537
- ☐ 11: [Muller WD, Karamfilov T, Kahlert H, Stuwe HT, Fahlbusch B, Cromwell O, Fiebig H, Jager L.](#) Related Articles
Mapping of T-cell epitopes of Phl p 5: evidence for crossreacting and non-crossreacting T-cell epitopes within Phl p 5 isoallergens.
Clin Exp Allergy. 1998 Dec;28(12):1538-48.
PMID: 10024226
- ☐ 12: [Mizumoto K, Kimura S, Abe Y, Uehara M, Katagiri M.](#) Related Articles
[Analysis of T cell epitopes on birch pollen allergen].
Hokkaido Igaku Zasshi. 1997 Jan;72(1):59-67. Japanese.
PMID: 9086363
- ☐ 13: [Hashiguchi S, Hino K, Taniguchi Y, Kurimoto M, Fukuda K, Ohyama M, Fujiyoshi T, Sonoda S, Nishimura Y, Yamada G, Sugimura K.](#) Related Articles
Immunodominance of seven regions of a major allergen, Cry j 2, of Japanese cedar pollen for T-cell immunity.
Allergy. 1996 Sep;51(9):621-32.
PMID: 8899114
- ☐ 14: [Mori T, Yokoyama M, Komiyama N, Okano M, Kino K.](#) Related Articles, Protein
Purification, identification, and cDNA cloning of Cha o 2, the second major allergen of Japanese cypress pollen.
Biochem Biophys Res Commun. 1999 Sep 16;263(1):166-71.
PMID: 10486272
- ☐ 15: [Burton MD, Blaher B, Suphioglu C, O'Hehir RE, Carbone FR, Rolland JM.](#) Related Articles
T-cell receptor contact and MHC binding residues of a major rye grass pollen allergen T-cell epitope.
J Allergy Clin Immunol. 1999 Feb;103(2 Pt 1):255-61.

PMID: 9949316

- ☐ 16: [Hashimoto M, Kobayashi T, Nigi H, Saito S, Nakayama I, Narita T, Iwata M, Yasueda H, Taniguchi Y, Kurimoto M, Inouye S, Sakaguchi M.](#) [Related Articles](#)
 Responses of monkeys with pollinosis to two major allergens of Japanese cedar pollen.
 Int Arch Allergy Immunol. 1997 Jan;112(1):88-92.
 PMID: 8980469

-  17: [Ikagawa S, Matsushita S, Chen YZ, Ishikawa T, Nishimura Y.](#) [Related Articles](#)
 Single amino acid substitutions on a Japanese cedar pollen allergen (Cry j 1)-derived peptide induced alterations in human T cell responses and T cell receptor antagonism.
 J Allergy Clin Immunol. 1996 Jan;97(1 Pt 1):53-64.
 PMID: 8568138


- ☐ 18: [Imai T, Nohara O, Katayama N, Endo T, Inoue S, Sakaguchi M, Taniguchi Y, Nakagawa T, Saito S.](#) [Related Articles](#)
 [Evaluation of efficacy for Japanese cedar pollen-specific immunotherapy by lymphocyte proliferation test].
 Arerugi. 1996 Nov;45(11):1172-80. Japanese.
 PMID: 8990529

- ☐ 19: [Nohara O, Imai T, Endo T, Saneyoshi K, Ohmori T, Saito S, Taniguchi Y, Fujimaki H, Moriyama H.](#) [Related Articles](#)
 [Relationship between the morphological changes of Japanese cedar (Cryptomeria japonica) pollen grains and the release of major allergens from the pollen].
 Arerugi. 1997 Dec;46(12):1235-42. Japanese.
 PMID: 9503683

- ☐ 20: [Costa MA, Duro G, Izzo V, Colombo P, Mirisola MG, Locorotondo G, Cocchiara R, Geraci D.](#) [Related Articles](#)
 The IgE-binding epitopes of rPar j 2, a major allergen of Parietaria judaica pollen, are heterogeneously recognized among allergic subjects.
 Allergy. 2000 Mar;55(3):246-50.
 PMID: 10753015

Display	Summary	<input type="checkbox"/> Save	Text	Order	Add to Clipboard
Show	20	<input type="checkbox"/> Items 1-20 of 216	Page 1 of 11	Select page	1 2 3 4 5 6 7 8 9 10 >>

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History	Clipboard		

Entrez PubMed

☐ 1: *Allergy* 1996 Sep;51(9):621-32[Related Articles, Books, LinkOut](#)PubMed
Services**Immunodominance of seven regions of a major allergen, Cry j 2, of Japanese cedar pollen for T-cell immunity.****Hashiguchi S, Hino K, Taniguchi Y, Kurimoto M, Fukuda K, Ohyama M, Fujiyoshi T, Sonoda S, Nishimura Y, Yamada G, Sugimura K**

Department of Molecular Biology, Faculty of Engineering, Kagoshima University, Japan.

Related
Resources

The immunodominant regions of the Japanese cedar pollen allergen Cry j 2 for T-cell immunity were determined with whole peripheral blood lymphocytes (PBL) derived from seven allergic patients and three nonallergic subjects. Cry j 2-stimulated T-cell proliferation was inhibited by anti-HLA-DR, but not by anti-HLA-DQ antibody, indicating that the responding T cells recognized the allergen peptides associated with HLA-DR molecules. It was found that seven regions of Cry j 2, i.e., regions corresponding to amino acid numbers 1-26, 70-84, 151-167, 187-203, 252-279, 283-314, and 345-362, were immunodominant for T-cell proliferation. Thus, Cry j 2 bears a limited number of immunodominant regions despite polymorphic features of HLA-DR in the immune system. This suggests the possibility of molecularly designing Cry j 2 antagonists that could downregulate allergic reactions to Japanese cedar pollen.

PMID: 8899114

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

[Write to the Help Desk](#)[NCBI](#) | [NLN](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)